

# Variational Principles

Hasan Baig


Easter 2021

## Contents

<b>0</b>	<b>Motivation</b>	<b>2</b>
<b>1</b>	<b>Calculus for Functions of <math>\mathbb{R}^n</math></b>	<b>5</b>
1.1	Constraints and Lagrange Multipliers . . . . .	7
<b>2</b>	<b>Euler-Lagrange Equations</b>	<b>9</b>
2.1	First Integrals of the E-L equation . . . . .	11
2.2	Fermat's Principle . . . . .	16
<b>3</b>	<b>Extensions of the Euler-Lagrange Equations</b>	<b>18</b>
3.1	Euler-Lagrange Equations with Constraints . . . . .	18
3.2	Several dependent variables . . . . .	20
3.3	Several Independent Variables . . . . .	22
3.4	Higher Derivatives . . . . .	25
<b>4</b>	<b>Least Action Principle and Noether's Theorem</b>	<b>26</b>
4.1	Noether's Theorem . . . . .	28
<b>5</b>	<b>Convex Functions</b>	<b>29</b>
5.1	Conditions for Convexity . . . . .	31
<b>6</b>	<b>Legendre Transform</b>	<b>32</b>
6.1	Applications to Thermodynamics . . . . .	34
<b>7</b>	<b>Hamilton's Equations</b>	<b>34</b>
<b>8</b>	<b>The Second Variation</b>	<b>36</b>
8.1	Associated Eigenvalue Problem . . . . .	38
8.2	The Jacobi Condition . . . . .	39

## 0 Motivation

**Example** (The Brachistochrone Problem). A particle slides on a wire, under influence of gravity between two fixed points  $A$ ,  $B$ . Which shape of the wire gives the shortest travel time, starting from rest?



Johann Bernoulli proposed the problem of finding the optimal shape, in 1696. Travel time:

$$T = \int dt = \int_A^B \frac{dl}{v(x,y)}$$

$$K.E. + V = \text{const. (energy conservation)}$$


$$\frac{1}{2}mv^2 + mgy = mgy_1 = 0 \qquad v = \sqrt{2g\sqrt{-y}}$$

Minimise

$$T[y] = \frac{1}{\sqrt{2g}} \int_0^{x_2} \frac{\sqrt{1+(y')^2}}{\sqrt{-y}} dx$$

subject to  $y(0) = 0$ ,  $y(x_2) = y_2$ .

**Example** (Geodesic). Finding the shortest path  $\gamma$  between 2 points on a surface  $\Sigma$  (if one exists). Take  $\Sigma = \mathbb{R}^2$  (a plane, Pythagorean theorem holds).



Distance along  $\gamma$ :

$$D[y] = \int_A^B dl = \int_{x_1}^{x_2} \sqrt{1 + (y')^2} dx$$

Seek to minimise  $D$  by varying  $\gamma$ .


**Remark.** Generally, we are trying to minimise (maximise)

$$F[y] = \int_{x_1}^{x_2} f(x, y(x), y'(x)) dx \tag{0.1}$$

among all functions s.t.  $y(x_1) = y_1, y(x_2) = y_2$ .

(0.1) is a **functional** (a function on the space of functions)

Functions map numbers to numbers. Functionals map functions to numbers e.g.



$y(x) \rightarrow$  area under the graph

$$f(x, y, y') = y$$

• curve  $\rightarrow$  length

$$f(x, y, y') = \sqrt{1 + (y')^2}$$

Calculus of variations is finding extrema (min/max/stable) of functions on spaces of functions.

**Notation.**  $C(\mathbb{R})$  is the space of continuous functions on  $\mathbb{R}$   
 $C^k(\mathbb{R})$  is the space of continuous functions on  $\mathbb{R}$  with continuous  $k$ -th derivatives  
 $C^k_{\alpha,\beta}(\mathbb{R})$  is the space of continuous functions on  $\mathbb{R}$  with continuous  $k$ -th derivatives s.t.  $f(\alpha) = f(\beta)$


**Warning.** NEED to specify the function space beforehand (a branch of Functional Analysis – Part III – analysis on the space of functions)

Variational Principles are principles in nature where the laws follow from extremising Functionals

**Example** (Fermat’s Principle). “Light between two points travels along paths which require least time.”

**Example** (Principle of least action).  $T =$  kinetic energy (e.g.  $m|\dot{\mathbf{x}}|^2/2$ )  
 $V =$  potential energy (e.g.  $V(\mathbf{x}), \mathbf{x} \in \mathbb{R}^n$ )

$$S[\gamma] = \int_{t_1}^{t_2} (T - V) dt$$



“Action is minimised along paths of motion”

**Moral.** Leibnitz’s take: We live in “the best of all possible worlds”.

Science → Theology.

Feynman’s take: “This is wrong. In quantum theory, the motion takes place along all possible paths with different probabilities.” (see Part III QFT)

In this course

- We consider necessary conditions of extremum of (0.1). Euler-Lagrange equation.
- Lots of examples (geometry, physics, problems with constraints – e.g. maximise area given a fixed length of perimeter)
- Second variation: some sufficient conditions for min/ max

Books:

- (i) Gelfand- Fomin 'Calculus of Variations.'
- (ii) DAMTP notes online (e.g. P. Townsend)

**Note.** Lectures have a different order but similar content to (ii).

## 1 Calculus for Functions of $\mathbb{R}^n$

In this section,  $f \in C^2(\mathbb{R}^n)$ ,  $f: \mathbb{R}^n \rightarrow \mathbb{R}$ , continuous 2<sup>nd</sup> partial derivatives.

**Definition.** The position  $\mathbf{a} \in \mathbb{R}^n$  is **stationary** if

$$\nabla f(\mathbf{a}) = (\partial_1 f, \dots, \partial_n f) |_{\mathbf{x}=\mathbf{a}} = 0, \text{ where } \partial_i f = \frac{\partial f}{\partial x_i}$$

**Method.** Expanding near  $\mathbf{x} = \mathbf{a}$

$$f(\mathbf{x}) \stackrel{+}{\neq} f(\mathbf{a}) = \underbrace{(\mathbf{x} - \mathbf{a}) \cdot \partial f |_{\mathbf{a}}}_{0, \text{ as stationary}} + \frac{1}{2} (x_i - a_i)(x_j - a_j) \partial_{ij}^2 f |_{\mathbf{a}} + O(|\mathbf{x} - \mathbf{a}|^2)$$

using the summation convention. The **Hessian** matrix is

$$H_{ij} = \partial_i \partial_j f = H_{ji}$$

We shift the origin to set  $\mathbf{a} = \mathbf{0}$ , and diagonalise  $H(\mathbf{0})$  by an orthogonal transformation:

$$H' = R^T H(\mathbf{0}) R = \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}$$

$$f(\mathbf{x}') - f(\mathbf{0}) = \frac{1}{2} \sum \lambda_i (x'_i)^2 + O(|\mathbf{x}'|^2), \quad \mathbf{x}' = R^T \mathbf{x}$$

- (i) If all  $\lambda_i > 0$ ,  $f(\mathbf{x}') > f(\mathbf{0})$  in all directions so we have a local minimum
- (ii) If all  $\lambda_i < 0$ , then we have a local maximum
- (iii) If some  $\lambda_i > 0$ , and some  $\lambda_i < 0$ , then  $f$  increases in some directions and decreases in others. We have a **saddle point** in this case.
- (iv) If some  $\lambda_i = 0$ , then we need to consider higher order derivatives in Taylor's expansion. If some are +ve and some -ve, then we know its a saddle point even if there are 0s.

**Method.** Special case  $n = 2$ :

$$\det(H) = \lambda_1 \lambda_2, \quad \text{tr}(H) = \lambda_1 + \lambda_2$$

- $\det > 0$ ,  $\text{tr} > 0$  gives local minimum
- $\det > 0$ ,  $\text{tr} < 0$  gives local maximum
- $\det < 0$  gives saddle point
- $\det = 0$  requires us to look at 3<sup>rd</sup>/ higher derivatives

max on boundary.



**Remarks.**

- (i) For  $f : D \rightarrow \mathbb{R}$  (domain) we can have local minimum, local maximum or global minimum
- (ii) For  $f$  harmonic,  $f_{xx} + f_{yy} = 0$ ,  $D \subseteq \mathbb{R}^2$  gives  $\text{tr}(H) = 0$  so our turning point is a saddle point and the min/ max is on the boundary

**Example.**

$$f(x, y) = x^3 + y^3 - 3xy$$

$$\nabla f = (3x^2 - 3y, 3y^2 - 3x) = (0, 0)$$

for critical points.


$$x^2 - y = 0, y^2 = 0 \implies y^4 = y \implies \begin{cases} y = 0, x = 0 \\ y = 1, x = 1 \end{cases}$$

Stationary points  $(0, 0)$  and  $(1, 1)$

$$H = \begin{bmatrix} 6x & -3 \\ -3 & 6y \end{bmatrix}$$

$(0, 0)$  has  $\det H = -9 < 0$ , saddle point  $f = 0$ .


$(1, 1)$  has  $\det H = 27 > 0$ ,  $\text{tr}(H) = 12 > 0$  so is local minimum with  $f = -1$



For  $(0, 0)$ , near  $f = 0$ ,  $f \simeq -3xy$  which decreases on the line  $y = x$  but increases on  $y = -x$ .  
This function has no global min/ max

## 1.1 Constraints and Lagrange Multipliers

**Example.** Find the circle centered at  $(0,0)$ , with smallest radius, which intersects the parabola  $y = x^2 - 1$



Two approaches:

(i) Direct method. Solve the constraints

$$f = x^2 + y^2 = x^2 + (\underbrace{x^2 - 1}_{\mathcal{C}})^2 = \underbrace{x^4 - x^2 + 1}_{f(x)}$$

We have

$$\partial_x f = 0 \iff 4x^3 - 2x = 0$$

Giving two solutions

- $x = \pm 1/\sqrt{2}$ ,  $y = -1/2$ , radius  $\sqrt{3}/2$
- $x = 0$ ,  $y = -1$ , radius 1

**Example.** (ii) Lagrange Multipliers. Define new function  $h(x, y, \lambda) = f(x, y) - \lambda g(x, y)$  with  $g(x, y) = 0$  the constraint.  $\lambda =$  Lagrange multiplier.

$$h = x^2 + y^2 - \lambda(y - x^2 + 1)$$

Extremising over 3 variables with no constraints:

$$\frac{\partial h}{\partial x} = 2x + 2\lambda x = 0$$

$$\frac{\partial h}{\partial y} = 2y - \lambda = 0$$

$$\frac{\partial h}{\partial \lambda} = y - x^2 + 1 = 0$$

The first two give:

$$2x + 4xy = 0 \implies x = 0 \text{ or } y = -\frac{1}{2}$$


Subbing these in the final equation gives solutions:

$$(x, y) = (0, 1) \text{ or } \left(\pm \frac{1}{\sqrt{2}}, -\frac{1}{2}\right)$$

$$(0, 1) \rightarrow f = 1 \text{ so } (\lambda = 2)$$

$$\left(\pm \frac{1}{\sqrt{2}}, -\frac{1}{2}\right) \rightarrow f = \frac{3}{4}, \lambda = -1$$

**Moral.** Why does it work (geometry):



At the extrema,  $\nabla f \parallel \nabla g$ , so

$$\nabla f = \lambda \nabla g \text{ i.e. } \nabla(f - \lambda g) = 0$$

Extremum of  $h = f - \lambda g$



**Method.** For multiple constraints, extremise  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ , subject to  $g_\alpha(\mathbf{x}) = 0$

$$g_\alpha : \mathbb{R}^n \rightarrow \mathbb{R} \quad \alpha = 1, \dots, k$$

$$h(x_1, \dots, x_n, \lambda_1, \dots, \lambda_k) = f - \sum_{\alpha=1}^k \lambda_\alpha g_\alpha$$

We have  $n + k$  variables,  $k$  Lagrange Multipliers

$$\frac{\partial h}{\partial x_i} = 0, \quad \frac{\partial h}{\partial \lambda_\alpha} = 0$$


Eliminate  $\lambda_\alpha$  and solve for  $\mathbf{x}$

This method works also if constraints can't be eliminated

## 2 Euler-Lagrange Equations

**Method.** Our task is to extremise functional (0.1)

$$F[y] = \int_{\alpha}^{\beta} f(x, y, y') dx$$



$f$  given, depends on  $y$  with fixed ends.

Consider a small perturbation  $y \rightarrow y + \epsilon\eta(x)$  in (2.1)

Compute  $F[y + \epsilon\eta]$ ,  $\eta(\alpha) = \eta(\beta) = 0$ .

We will need the lemma below

**Lemma.** If  $g : [\alpha, \beta] \rightarrow \mathbb{R}$  is continuous on  $[\alpha, \beta]$ , and

$$\int_{\alpha}^{\beta} g(x)\eta(x) dx = 0 \text{ for all } \eta \text{ continuous on } [\alpha, \beta], \text{ s.t. } \eta(\alpha) = \eta(\beta) = 0$$

Then  $g(x) \equiv 0, \forall x \in [\alpha, \beta]$

**Proof.** We show  $\exists \bar{x} \in (\alpha, \beta)$  s.t.  $g(\bar{x}) \neq 0$ . Suppose  $g(\bar{x}) > 0$ . Then  $\exists$  interval  $[x_1, x_2] \subseteq [\alpha, \beta]$  s.t.  $g(x) > c$  on  $[x_1, x_2]$  for some  $c > 0$ . Set

$$\eta(x) = \begin{cases} (x - x_1)(x_2 - x) & x \in [x_1, x_2] \\ 0 & x \notin [x_1, x_2] \end{cases} \quad (2.2)$$

$$\int_{\alpha}^{\beta} g(x)\eta(x) dx > c \int_{x_1}^{x_2} (x - x_1)(x_2 - x) dx > 0$$

**Remark.**  $\eta$  given by (2.2) is a bump function. A  $C^k$  bump function:

$$\eta = \begin{cases} ((x - x_1)(x_2 - x))^{k+1} & x \in [x_1, x_2] \\ 0 & x \notin [x_1, x_2] \end{cases}$$

**Method.** Back to (2.1):

$$\begin{aligned} F[y + \varepsilon\eta] &= \int_{\alpha}^{\beta} f(x, y + \varepsilon\eta, y' + \varepsilon\eta') dx \\ &= F[y] + \varepsilon \int_{\alpha}^{\beta} \left( \frac{\partial f}{\partial y} \eta + \frac{\partial f}{\partial y'} \eta' \right) dx + \underbrace{O(\varepsilon^2)}_{\text{return in section 8}} \\ &= F[y] + O(\varepsilon^2) \text{ at extremum, i.e. } \left. \frac{dF}{d\varepsilon} \right|_{\varepsilon=0} = 0 \end{aligned}$$

Integrating the  $\varepsilon$ -term by parts

$$\begin{aligned} 0 &= \int_{\alpha}^{\beta} \left\{ \frac{\partial f}{\partial y} \eta - \frac{d}{dx} \left( \frac{\partial f}{\partial y'} \eta \right) \right\} dx + \underbrace{\left[ \frac{\partial f}{\partial y'} \eta \right]_{\alpha}^{\beta}}_{0 \text{ as } \eta(\alpha) = \eta(\beta) = 0} \\ &= \int_{\alpha}^{\beta} \underbrace{\left( \frac{\partial f}{\partial y} - \frac{d}{dx} \left( \frac{\partial f}{\partial y'} \right) \right)}_{=g} \eta dx \end{aligned}$$

$\int \left( \frac{\partial f}{\partial y} \eta + \frac{\partial f}{\partial y'} \eta' \right) dx = 0, \int \frac{\partial f}{\partial y'} \eta' dx = \left[ \frac{\partial f}{\partial y'} \eta \right] - \int \eta \frac{d}{dx} \left( \frac{\partial f}{\partial y'} \right) dx.$

Applying the Lemma with  $g$  as above, we must have  $g \equiv 0$ .

**Equation.** We have proved a necessary condition for an extremum is:

$$\frac{d}{dx} \left( \frac{\partial f}{\partial y'} \right) - \frac{\partial f}{\partial y} = 0 \quad (2.3)$$

This is called the Euler-Lagrange equation

**Remarks.**

- (2.3) is a 2nd order ODE for  $y(x)$  with boundary conditions  $y(\alpha) = y_1, y(\beta) = y_2$
- Notation: the LHS of (2.3) denoted by  $\frac{\partial F}{\partial y(x)}$  is called the functional derivatives
- Some books (e.g. Towsend's notes) use  $\delta y = \varepsilon \eta(x)$

$$F[y + \delta y] = F[y] + \delta F[y]$$

where

$$\delta F = \int_{\alpha}^{\beta} \left[ \frac{\partial F(y)}{\partial y(x)} \delta y(x) \right] dx$$

- Other boundary conditions are possible e.g.  $\frac{\partial f}{\partial y'}|_{\alpha, \beta} = 0$
- Be careful with derivatives, e.g.  $\frac{\partial f}{\partial y}$  means  $(\frac{\partial f}{\partial y})_{x, y'}$  ( $x, y, y'$  independent in  $f$ )

$$\frac{dh}{dx} = \frac{\partial h}{\partial x} + \frac{\partial h}{\partial y} y' + \frac{\partial h}{\partial y'} y''$$

$$\frac{d}{dx} = \delta_x + y' \delta_y + y'' \delta_{y'}$$

is the total derivative.

**Example.**

$$\begin{aligned} f(x, y, y') &= x \cdot ((y')^2 - y^2) \\ \delta_x f &= (y')^2 - y^2 \quad \delta_y f = -2xy \quad \delta_{y'} f = 2xy' \\ \frac{df}{dx} &= (y')^2 - y^2 - 2xyy' + 2xy'y'' \end{aligned}$$

## 2.1 First Integrals of the E-L equation


In some cases (2.3) (2nd order ODE) can be integrated once to a 1st order ODE "first integral".

- (i)  $f$  does not explicitly depend on  $y$ ,  $\frac{df}{dy} = 0$

$$\frac{\partial f}{\partial y} = 0$$

$$(2.3) \rightarrow \frac{d}{dx} \left( \frac{\partial f}{\partial y'} \right) = 0 \implies \frac{\partial f}{\partial y'} = \text{constant}$$

**Examples.** Geodesics on the Euclidean plane



$$F[y] = \int_{\alpha}^{\beta} \sqrt{dx^2 + dy^2} = \int_{\alpha}^{\beta} \underbrace{\sqrt{1 + (y')^2}}_{f(y')} dx$$

$$\frac{\partial f}{\partial y} = 0 \implies \frac{y'}{\sqrt{1 + (y')^2}} = \text{const.} = \frac{\partial f}{\partial y'}$$

So


$$y' = m$$

for some constant  $m$ , and so

$$y = mx + c$$

straight line

**Example.** Geodesics on a sphere  $S^2 \subset \mathbb{R}^3$



$$x = \sin \theta \sin \phi \quad 0 \leq \theta \leq \pi$$

$$y = \sin \theta \cos \phi \quad 0 \leq \phi \leq 2\pi$$

$$z = \cos \theta$$

$$ds^2 = dx^2 + dy^2 + dz^2 = d\theta^2 + \sin^2 \theta d\phi^2$$

Parametrise as  $\phi = \phi(\theta)$

$$ds = \sqrt{1 + \sin^2 \theta (\phi')^2} d\theta$$

$$F[\phi] = \int_{\theta_1=\alpha}^{\theta_2=\beta} \sqrt{1 + \sin^2 \theta \cdot (\phi')^2} d\theta$$

$$\frac{\partial f}{\partial \phi} = 0 \implies \frac{\partial f}{\partial \phi'} = \kappa \text{ (constant)}$$

first integral.

$$\frac{\sin^2 \theta \cdot \phi'}{\sqrt{1 + \sin^2 \theta \cdot (\phi')^2}} = \kappa$$

**Example** (continued). Squaring to solve for  $(\phi')^2$


$$(\phi')^2 = \frac{\kappa^2}{\sin^2 \theta \cdot (\sin^2 \theta - \kappa^2)}$$

$$\phi = \pm \int \frac{\kappa d\theta}{\sin \theta \cdot \sqrt{\sin^2 \theta - \kappa^2}}$$

Two solutions, each going one way around the sphere. Using substitution  $\cot(\theta) = u$

$$\pm \frac{\sqrt{1 - \kappa^2}}{\kappa} \cos(\phi - \phi_0) = \cot \theta$$

for  $\phi_0 = \text{const.}$  Great circle (plane through  $\odot$  intersecting  $S^2$ ).



(Geodesics are segments of great circles)

(ii) Consider, for general  $f(x, y, y')$


$$\begin{aligned} \frac{d}{dx} \left( f - y' \frac{\partial f}{\partial y'} \right) &= \frac{\partial f}{\partial x} + y' \frac{\partial f}{\partial y} + y'' \frac{\partial f}{\partial y'} - \cancel{y'' \frac{\partial f}{\partial y'}} - y' \frac{d}{dx} \left( \frac{\partial f}{\partial y'} \right) \\ &= y' \underbrace{\left( \frac{\partial f}{\partial y} - \frac{d}{dx} \frac{\partial f}{\partial y'} \right)}_{=0} + \frac{\partial f}{\partial x} \end{aligned}$$

↪ Euler Lagrange

If  $f$  does not explicitly depend on  $x$ , i.e.  $\frac{\partial f}{\partial x} = 0$  then

$$f - y' \frac{\partial f}{\partial y'} = \text{const.} \quad (2.5)$$

Example (Brachistochrone).



Going back to section 0,

$$F[y] = \frac{1}{\sqrt{2g}} \int_0^\beta \underbrace{\frac{\sqrt{1+(y')^2}}{\sqrt{-y}}}_{f(y,y')} dx$$

$\frac{\partial f}{\partial x} = 0$  so use (2.5)

$$\frac{\sqrt{1+(y')^2}}{\sqrt{-y}} - y' \frac{y'}{\sqrt{1+(y')^2}\sqrt{-y}} = K \quad \text{K includes g term}$$

$$\frac{1}{\sqrt{1+(y')^2}} = K\sqrt{-y} \implies y' = \pm \frac{\sqrt{1+K^2 y^2}}{K\sqrt{-y}}$$

$$x = \pm K \int \frac{\sqrt{-y}}{\sqrt{1+K^2 y^2}} dy$$

Set

$$y = -\frac{1}{K^2} \sin^2 \frac{\theta}{2} \quad dy = -\frac{1}{K^2} \sin\left(\frac{\theta}{2}\right) \cos \frac{\theta}{2}$$

$$x = \pm K \int (-1) \frac{1}{K^2} \frac{\sin^2(\frac{\theta}{2}) \cos(\frac{\theta}{2})}{\sqrt{1 - \sin^2(\frac{\theta}{2})}} d\theta$$


$$= \mp \frac{1}{2K^2} \int (1 - \cos \theta) d\theta = \mp \frac{1}{2K^2} (\theta - \sin \theta) + C$$

Initial condition  $(0,0) \rightarrow \theta_0 = 0 \rightarrow C = 0$ , take positive root

$$x = \frac{\theta - \sin \theta}{2K^2}$$

$$y = -\frac{1}{K^2} \sin^2 \frac{\theta}{2}$$

**Example** (continued).



Parametrised equation of a cycloid. Brachistochrome = cycloid.  
The curve traced by a point on the rim of a wheel, as the wheel rolls along a straight line (Galileo)

## 2.2 Fermat's Principle

Light/sound travels along paths between two points which requires least time. Rays are represented by path  $y = y(x)$ . Speed of light  $c(x, y)$


$$F[y] = \int \frac{dl}{c} = \int_{\alpha}^{\beta} \frac{\sqrt{1 + (y')^2}}{c(x, y)} dx$$

assume  $c = c(x) \rightarrow \frac{\partial f}{\partial y} = 0$  so (2.4) gives

$$\frac{\partial f}{\partial y'} = \text{const.}$$

$$\frac{y'}{\sqrt{1 + (y')^2}c(x)} = \text{const.}$$






Ray launched at  $\theta_1$ ,  $\tan \theta = y'$


$$\frac{\sin \theta_1}{c(x_1)} = \frac{\sin \theta}{c(x)}$$

(2.6)


Snell's law.



$c$  increasing



$c$  decreasing



### 3 Extensions of the Euler-Lagrange Equations

#### 3.1 Euler-Lagrange Equations with Constraints

Extremise

$$F[y] = \int_{\alpha}^{\beta} f(x, y, y') dx$$

subject to

$$G[y] = \int_{\alpha}^{\beta} g(x, y, y') dx = K \text{ (constant)}$$


Lagrange multiplier, extremise

$$\Phi[y; \lambda] = F[y] - \lambda G[y]$$


replace  $f$  in E-L by  $f - \lambda g$

$$\frac{d}{dx} \left( \frac{\partial}{\partial y'} (f - \lambda g) \right) - \frac{\partial}{\partial y} (f - \lambda g) = 0 \tag{3.1}$$

**Example.** Dido problem (a.k.a. isoperimetric problem). What simple and closed plane curve of fixed length  $L$  maximises the enclosed area?




A simple curve doesn't intersect itself and it on a simply connected domain.



Not simple

Not convex  $\rightarrow$  convex with the same perimeter

Assume convexity.



$$dA = y(x) \Big|_{x_1}^{x_2} dx$$

$x$  monotonically increases from  $\alpha \rightarrow \beta$  and decreases from  $\beta \rightarrow \alpha$ . Given  $x, \exists (y_1, y_2)$  on the curve with

$$y_1(x) = y_2(x), \quad y_2 > y_1, \quad dA = y(x) \Big|_{x_1}^{x_2} \cdot dx$$

$$A[y] = \int_{\alpha}^{\beta} (y_2(x) - y_1(x)) dx = \oint_C y(x) dy$$

Constraint

$$L[y] = \oint_C dl = \oint_C \sqrt{1 + (y')^2} dx = L$$

$$h K = y \bar{\theta} \lambda \sqrt{1 + (y')^2}$$

(Note: do not worry about the boundary term in the derivation of the E-L, as  $C$  has no boundary)  
 $\frac{\partial h}{\partial x} = 0$  so we use (2.5)

$$K = \text{const} = h - y' \frac{\partial h}{\partial y'} = y - \lambda \sqrt{1 + (y')^2} + y' \lambda \frac{y'}{\sqrt{1 + (y')^2}}$$

$$\implies K = y - \frac{\lambda}{\sqrt{1 + (y')^2}} \implies (y')^2 = \frac{\lambda^2}{(y - k)^2} - 1$$

solution  $(x - x_0)^2 + (y - y_0)^2 = \lambda^2$  (circle of radius  $\lambda$ )

$$2\pi\lambda = L \implies \lambda = \frac{L}{2\pi}$$

**Example.** The Sturm-Liouville problem.

$\rho(x) > 0$  for  $x \in [\alpha, \beta]$ ,  $\sigma = \sigma(x)$

$$F[y] = \int_{\alpha}^{\beta} [\rho \cdot (y')^2 + \sigma y^2] dx \quad G[y] = \int_{\alpha}^{\beta} y^2 dx$$

Minimise  $f$  subject to  $G = 1$  (fixed ends) *(y(α), y(β) fixed)*

$$\Phi[y; \lambda] = F[y] - \lambda(G[y] - 1)$$

$$h = \rho \cdot (y')^2 + \sigma \cdot y^2 - \lambda \left( y^2 - \frac{1}{\beta - \alpha} \right) \quad \int_{\alpha}^{\beta} \frac{1}{\beta - \alpha} = 1.$$

$$\frac{\partial h}{\partial y'} = 2\rho y' \quad \frac{\partial h}{\partial y} = 2\sigma y - 2\lambda y$$

$$-\frac{d}{dx} (\rho \cdot y') + \sigma \cdot y = \lambda y \quad (3.2)$$

$\underbrace{\hspace{10em}}_{\mathcal{L}(y)}$

$\mathcal{L}$  is the Sturm-Liouville operator. (3.2) is an eigenvalue problem. e.g. if  $\rho = 1$ ,  $\sigma(x) =$  'potential' in time-independent Schrödinger equation (IB Quantum Mechanics).

If  $\sigma > 0$ , then  $F[y] > 0$ . Positive minimum equal to the lowest eigenvalue

**Proof.** (3.2)  $\times y$  and integrate  $\int_{\beta}^{\alpha}$  by parts

$$F[y] - \underbrace{[y \cdot y' \rho]_{\beta}^{\alpha}}_0 = \underbrace{G[y]}_1 \cdot \lambda$$

Lowest eigenvalue is the minimum of  $F[y]/G[y]$  *Assume  $y(\alpha) = y(\beta) = 0$ .*

### 3.2 Several dependent variables

$$\mathbf{y}(x) = (y_1(x), y_2(x), \dots, y_n(x))$$

$$F[\mathbf{y}] = \int_{\alpha}^{\beta} f(x, y_1, \dots, y_n, y'_1, \dots, y'_n) dx$$

$$y_i \rightarrow y_i(x) + \varepsilon \eta_i(x) \quad i = 1, \dots, n \quad \eta_i(\alpha) = \eta_i(\beta) = 0$$

Following the derivation of the E-L equation: *(Assume  $\mathbf{y}$  is optimal)*

$$F[\mathbf{y} + \varepsilon \boldsymbol{\eta}] - F[\mathbf{y}] = \int_{\alpha}^{\beta} \sum_{i=1}^n \eta_i \left( \frac{d}{dx} \left( \frac{\partial f}{\partial y'_i} \right) - \frac{\partial f}{\partial y_i} \right) dx + \text{boundary term} + O(\varepsilon^2)$$

Use Lemma *(Letting  $\eta_i = \delta_{ij} \eta$ )*

$$\frac{d}{dx} \left( \frac{\partial f}{\partial y'_i} \right) = \frac{\partial f}{\partial y_i} \quad (3.3)$$

A system of  $n$  2nd order ODEs.

First integrals of 3.3

- If  $\frac{\partial f}{\partial y_j} = 0$  for some  $1 \leq j \leq n$  then, by (3.3)  $\frac{\partial f}{\partial y'_j} = \text{const.}$
- If  $\frac{\partial f}{\partial x} = 0$ , then  $f - \sum_i y'_i \frac{\partial f}{\partial y'_i} = \text{const.}$

**Example.** Geodesics on surfaces

$\Sigma \subset \mathbb{R}^3$  (surface) given by

$$g(x, y, z) = 0$$

**Geodesic** = shortest path on the surface between  $A, B \in \Sigma$  (if one exists).  $t$  = parameter on the curve

$$A = \mathbf{x}(0)$$

$$B = \mathbf{x}(1) \quad \mathbf{x} = (x, y, z)$$

$$\Phi[\mathbf{x}, \lambda] = \int_0^1 \underbrace{\sqrt{\dot{x}^2 + \dot{y}^2 + \dot{z}^2} - \lambda(t) \cdot g(x, y, z)}_{h(x, y, z, \dot{x}, \dot{y}, \dot{z}, \lambda)} dt$$

Note: The Lagrange multiplier  $\lambda$  is now a function of  $t$  as we want the entire curve to lie on  $\Sigma$ . E-L equations with  $h$ .

- Variation w.r.t.  $\lambda$ :

$$\underbrace{\frac{d}{dt} \left( \frac{\partial h}{\partial \dot{z}} \right)}_0 - \frac{\partial h}{\partial \lambda} = 0 \implies g(x, y, z) = 0 \quad \forall t$$

- Variation w.r.t.  $x_i = (x, y, z)$

$$\frac{d}{dt} \left( \frac{\dot{x}_i}{\sqrt{\dot{x}^2 + \dot{y}^2 + \dot{z}^2}} \right) + \lambda \frac{\partial g}{\partial x_i} = 0 \quad i = 1, 2, 3$$

**We can use the two equations to eliminate  $\lambda$**

Alternatively, solve the constraint  $g = 0$ , as we did in example 2.2 ( $\Sigma = \text{sphere}$ )

### 3.3 Several Independent Variables

In general  $\phi: \mathbb{R}^n \rightarrow \mathbb{R}^m$ . In  $n > 1$ , E-L become PDEs. Assume that  $n = 3, m = 1$

$\phi$

$$F[\phi] = \iiint_D \underbrace{f(x, y, z, \phi, \phi_x, \phi_y, \phi_z)}_{\text{indep}}, dx dy dz$$

notation  $\phi_x = \frac{\partial \phi}{\partial x}$  etc. Volume integral over a domain  $D \subset \mathbb{R}^3$ . Assume  $\phi$  extremum, consider perturbations

$$\phi \rightarrow \phi(x, y, z) + \varepsilon \eta(x, y, z) \text{ s.t. } \eta = 0 \text{ on } \partial D$$

$$\begin{aligned} F[\phi + \varepsilon \eta] - F[\phi] &= \varepsilon \int_D \left( \eta \frac{\partial f}{\partial \phi} + \eta_x \frac{\partial f}{\partial \phi_x} + \eta_y \frac{\partial f}{\partial \phi_y} + \eta_z \frac{\partial f}{\partial \phi_z} \right) dx dy dz + O(\varepsilon^2) \\ &= \varepsilon \int_D \eta \frac{\partial f}{\partial \phi} + \nabla \cdot \left( \eta \left( \frac{\partial f}{\partial \phi_x}, \frac{\partial f}{\partial \phi_y}, \frac{\partial f}{\partial \phi_z} \right) \right) - \eta \nabla \cdot \left( \frac{\partial f}{\partial \phi_x}, \frac{\partial f}{\partial \phi_y}, \frac{\partial f}{\partial \phi_z} \right) dx dy dz + O(\varepsilon^2) \end{aligned}$$

Apply divergence theorem to first div term and use

$$\int_{\partial D} \eta \left( \frac{\partial f}{\partial \phi_x}, \frac{\partial f}{\partial \phi_y}, \frac{\partial f}{\partial \phi_z} \right) \cdot ds = 0$$

as  $\eta = 0$  on  $\partial D$

$$F[\phi + \varepsilon \eta] - F[\phi] = \varepsilon \int_D \eta \left( \frac{\partial f}{\partial \phi} - \nabla \cdot \left( \frac{\partial f}{\partial \phi_x}, \frac{\partial f}{\partial \phi_y}, \frac{\partial f}{\partial \phi_z} \right) \right) dx dy dz + O(\varepsilon^2)$$

E-L equation: single 2nd order PDE for one function  $\phi$

$$\frac{\partial f}{\partial \phi} - \sum_{i=1}^3 \frac{\partial}{\partial x_i} \left( \frac{\partial f}{\partial \phi_i} \right) = 0 \quad (3.4)$$

remains valid with  $3 \rightarrow n$

**Example.** Extremise ‘potential energy’  $n = 2$

$$F[\phi] = \iint_D \frac{1}{2} [\phi_x^2 + \phi_y^2] dx dy$$

$$\frac{\partial f}{\partial \phi} = 0 \quad \frac{\partial f}{\partial \phi_x} = \phi_x \quad \frac{\partial f}{\partial \phi_y} = \phi_y$$

$$(3.4) \rightarrow \frac{\partial \phi_x}{\partial x} + \frac{\partial \phi_y}{\partial y} = 0$$

i.e.

$$\phi_{xx} + \phi_{yy} = 0$$

(Laplace equation)

**Example.** Minimal surfaces. Minimise the area of  $\Sigma \subset \mathbb{R}^3$  subject to boundary conditions



e.g. soap forms

$$\Sigma = \{\mathbf{x} \in \mathbb{R}^3 : k(x, y, z) = 0\}$$

Assume (can do it by implicit function theorem) that we solved  $k = 0$  to give  $z = \phi(x, y)$

$$ds^2 = dx^2 + dy^2 + dz^2 \quad dz = \phi_x dx + \phi_y dy$$

$$ds^2 = (1 + \phi_x^2)dx^2 + (1 + \phi_y^2)dy^2 + 2\phi_x\phi_y dx dy$$

(IB geometry, this is called the 1st fundamental form or Riemannian metric)

$$ds^2 = \sum_{i,j=1}^2 g_{ij}(x, y) dx^i dx^j \quad x^1 = x, x^2 = y$$

$$g = \begin{bmatrix} 1 + \phi_x^2 & \phi_x\phi_y \\ \phi_x\phi_y & 1 + \phi_y^2 \end{bmatrix}$$

Area element  $\sqrt{\det g} dx dy$ . Area functional

$$A[\phi] = \int_D \sqrt{1 + \phi_x^2 + \phi_y^2} dx dy$$

Apply E-L (3.4) to  $h$

$$\frac{\partial h}{\partial \phi_x} = \frac{\phi_x}{\sqrt{1 + \phi_x^2 + \phi_y^2}} \quad \frac{\partial h}{\partial \phi_y} = \frac{\phi_y}{\sqrt{1 + \phi_x^2 + \phi_y^2}}$$

$$\partial_x \left( \frac{\phi_x}{\sqrt{1 + \phi_x^2 + \phi_y^2}} \right) + \partial_y \left( \frac{\phi_y}{\sqrt{1 + \phi_x^2 + \phi_y^2}} \right) = 0$$

Expand derivatives (exercise)

$$(1 + \phi_y^2)\phi_{xx} + (1 + \phi_x^2)\phi_{yy} - 2\phi_x\phi_y\phi_{xy} = 0 \quad (3.5)$$

The minimal surface equation. Assume circular symmetry

$$z = \phi(r) \quad r = \sqrt{x^2 + y^2}$$

$$\phi_x = \frac{dz}{dr} \frac{\partial r}{\partial x} = z' \frac{x}{r} \quad \phi_y = z' \frac{y}{r}$$

by calculating 2nd derivatives, we get from (3.5) the ODE

$$rz'' + z' + (z')^3 = 0$$


Set  $z' = w$  to get

$$\frac{1}{2}r \frac{dw^2}{dr} + w^2 + w^4 = 0$$

Solution

$$r = r_0 \cosh\left(\frac{z - z_0}{r_0}\right)$$

**Example** (continued). Catenoid: minimal surface of revolution (Euler 1744)




$r(L) = r(-L)$ . If  $L \neq 0$  then  $z_0 = 0$ . Set  $r = R$ , and divide by  $L$

$$\frac{R}{L} = \frac{r_0}{L} \cosh\left(\frac{L}{r_0}\right)$$


Set  $L = 1$ . Algebraic relation

$$R = r_0 \cosh(1/r_0)$$






**Example** (continued). If  $R \gg 1.5$ ,  $\exists 2$  minimal surfaces



unstable



stable

### 3.4 Higher Derivatives

**Equation.**

$$F[y] = \int_{\alpha}^{\beta} f(x, y, y', \dots, y^{(n)}) dx$$

Proceed as in section 2. Assume  $y$  exists,  $y \rightarrow t + \varepsilon\eta$  where

$$\eta = \eta' = \dots = \eta^{(n-1)} = 0 \text{ at } \alpha, \beta$$

$$F[y + \varepsilon\eta] - F[y] = \varepsilon \int_{\alpha}^{\beta} \left( \frac{\partial f}{\partial y} \eta + \frac{\partial f}{\partial y'} \eta' + \dots + \frac{\partial f}{\partial y^{(n)}} \eta^{(n)} \right) dx + O(\varepsilon^2)$$

integrate by parts n times  
 (blue arrows point from the text to the  $\eta^{(n)}$  term and from the  $\eta$  term to the  $\frac{\partial f}{\partial y}$  term)

Apply Lemma

$$\frac{\partial f}{\partial y} - \frac{d}{dx} \left( \frac{\partial f}{\partial y'} \right) + \frac{d^2}{dx^2} \left( \frac{\partial f}{\partial y''} \right) + \dots + (-1)^n \frac{d^n}{dx^n} \left( \frac{\partial f}{\partial y^{(n)}} \right) \quad (3.6)$$

Euler-Lagrange equation

**Example.** If  $n = 2$  and if  $\frac{\partial f}{\partial y} = 0$

$$(3.6) \rightarrow \frac{d}{dx} \left( \frac{\partial f}{\partial y'} - \frac{d}{dx} \frac{\partial f}{\partial y''} \right) = 0$$

so

$$\frac{\partial f}{\partial y'} - \frac{d}{dx} \frac{\partial f}{\partial y''} = \text{const.}$$

**Example.** Extremise  $F[y] = \int_0^1 (y'')^2 dx$  where  $y(0) = y'(0) = 0$  and  $y(1) = 0, y'(1) = 1$

$$\frac{d}{dx} (2y'') = \text{const.} \implies y''' = k \text{ const}$$

Impose boundary conditions to get  $y = x^3 - x^2$

**Note.** This is an absolute minimum.  $Y_0 = x^3 - x^2$

$$\eta(0) = \eta'(0) = \eta(1) = \eta'(1) = 0$$

(do not assume  $\eta$  small)

$$\begin{aligned} F[y_0 + \eta] - F[y_0] &= \int_0^1 (\eta'')^2 dx + 2 \cdot \int_0^1 (y_0'' \eta'') dx > 4 \int_0^1 (3x - 1) \eta'' dx \text{ if } \eta \neq 0 \forall x. \\ &= 4([-\eta']_0^1 + \int_0^1 \frac{d}{dx} (3x\eta') - \eta) dx \\ &= 4([3x\eta']_0^1 - 3\eta)_0^1 = 0 \end{aligned}$$

$y_0$  absolute minimiser of  $F$

## 4 Least Action Principle and Noether's Theorem

Particle  $\mathbb{R}^3, T = \text{kinetic energy}, V = \text{potential energy}.$

$$L(\mathbf{x}, \dot{\mathbf{x}}, t) = T - V \tag{4.1}$$

is the Lagrangian.  $t$  is the independent variable,  $\mathbf{x} = (x, y, z)$  are dependent variables. Action

$$S[\mathbf{x}] = \int_{t_1}^{t_2} L dt \tag{4.2}$$

Hamilton's principle (Least action principle, or principle of stationary action). The motion is such that  $S[\mathbf{x}]$  is stationary, i.e.  $L$  satisfies the E-L equations

**Example.**

$$T = \frac{1}{2} m |\dot{\mathbf{x}}|^2 \quad V = V(\mathbf{x})$$

Euler-Lagrange equations give

$$\begin{aligned} \frac{d}{dt} \left( \frac{\partial L}{\partial \dot{x}_i} \right) &= \frac{\partial L}{\partial x_i} \\ m \ddot{x}_i &= - \frac{\partial V}{\partial x_i} \text{ or } m \ddot{\mathbf{x}} = -\nabla V \end{aligned}$$

Newton's 2nd Law

**Example.** Central force in 2 dimensions:

$$L = \frac{1}{2}m(\dot{r}^2 + r^2\dot{\theta}^2) - V(r)$$

E-L

$$\frac{d}{dt} \left( \frac{\partial L}{\partial \dot{r}} \right) - \frac{\partial L}{\partial r} = 0$$

$$\frac{d}{dt} \left( \frac{\partial L}{\partial \dot{\theta}} \right) - \underbrace{\frac{\partial L}{\partial \theta}}_0 = 0$$

$$\implies \frac{\partial L}{\partial \dot{\theta}} = mr^2\dot{\theta} = \text{const.}$$

$\frac{\partial L}{\partial t} = 0$ , use (2.5)

$$\dot{r} \frac{\partial L}{\partial \dot{r}} + \dot{\theta} \frac{\partial L}{\partial \dot{\theta}} - L = \text{const}$$


$$\dot{r}mr\dot{r} + \dot{\theta}mr^2\dot{\theta} - \frac{1}{2}m\dot{r}^2 - \frac{1}{2}mr^2\dot{\theta}^2 + V(r) = \underbrace{\frac{1}{2}m\dot{r}^2 + \frac{1}{2}mr^2\dot{\theta}^2}_T + V(r) = E$$

which is constant. Conservation of total energy


**Example** (Configuration space, and general coordinates).

$N$  particles in  $\mathbb{R}^3$

$N = 3$



→



$$t \rightarrow \{q_i(t), \dot{q}_i(t), t\}$$

$q_i$  = generalised coordinates,  $i = 1, \dots, 3N$

$$\text{Langrangian } L = L(q_i, \dot{q}_i, t)$$

(Part II Classical Dynamics)

## 4.1 Noether's Theorem

$$F[\mathbf{y}] = \int_{\alpha}^{\beta} f(y_i, y_i', x) dx \quad i = 1, \dots, n$$

Suppose  $\exists$  a 1-parameter family of transformations  $y_i(x) \rightarrow Y_i(x, s)$  s.t.  $Y_i(x, 0) = y_i(x)$ . This is a continuous symmetry of a Lagrangian  $f$ , if

$$\frac{d}{ds} (f(Y_i(x, s), Y_i'(x, s), x)) = 0$$

**Theorem** (Noether's Theorem). Given a continuous symmetry  $Y_i(x, s)$  of  $f$ , the quantity

$$\sum_i \frac{\partial f}{\partial y_i'} \frac{\partial Y_i}{\partial s} \Big|_{s=0} \quad (4.3)$$

is a first integral of the E-L equation with  $Y_i(x, 0) = y_i(x) \forall i$

**Proof.**

$$\begin{aligned} 0 &= \frac{d}{ds} (f|_{s=0}) = \frac{\partial f}{\partial y_i} \frac{d y_i}{ds} \Big|_{s=0} + \frac{\partial f}{\partial y_i'} \frac{\partial Y_i'}{\partial s} \Big|_{s=0} \\ &= \left[ \frac{d}{dx} \left( \frac{\partial f}{\partial y_i'} \right) \frac{d Y_i}{ds} + \frac{\partial f}{\partial y_i'} \frac{d}{dx} \left( \frac{d Y_i}{ds} \right) \right] \Big|_{s=0} \quad \rightarrow \text{swap order of differentiation} \\ &= \frac{d}{dx} \left[ \frac{\partial f}{\partial y_i'} \frac{\partial Y_i}{\partial s} \right] \Big|_{s=0} = 0 \quad \leftarrow \text{by E-L} \end{aligned}$$

**Example.**

$$f = \frac{1}{2}(y')^2 + \frac{1}{2}(z')^2 - V(y - z), \quad \mathbf{y} = (y, z)$$

Lagrangian of a particle moving on a plane in a potential.

$$Y = y + s \quad Z = z + s \quad Y' = y' \quad Z' = z' \quad V(Y - Z) = V(y - z)$$

so

$$\begin{aligned} \frac{df}{ds} &= 0 \\ (4.3) \rightarrow \left( \frac{\partial f}{\partial y'} \frac{dY}{ds} + \frac{\partial f}{\partial z'} \frac{dZ}{ds} \right) &= y' + z' \end{aligned}$$

(conserved momentum in  $y + z$  direction)

**Example.** Back to example 4.2,  $\Theta = \theta + s, R = r$

$$\begin{aligned} \frac{dL}{ds} &= 0 \\ (4.3) \rightarrow \left( \frac{\partial L}{\partial \dot{\theta}} \frac{\partial \Theta}{\partial s} + \frac{\partial L}{\partial \dot{r}} \frac{\partial R}{\partial s} \right) \Big|_{s=0} &= mr^2 \dot{\theta} \end{aligned}$$


(conserved angular momentum). Isotropy of space gives rotational invariance of  $L$

## 5 Convex Functions


Going back to calculus on  $\mathbb{R}^n$ , a class of functions for which it is easy to classify stationary points

**Definition.** A set  $S \subset \mathbb{R}^n$  is **convex** if  $\forall \mathbf{x}, \mathbf{y} \in S$

$$(1-t)\mathbf{x} + t\mathbf{y} \in S \quad 0 \leq t \leq 1$$




convex



non-convex

**Definition.** A **graph of a function**  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is a surface

$$\{z - f(\mathbf{x}) = 0\} \text{ in } \mathbb{R}^{n+1}$$




A **chord** of  $f$  is a line segment in  $\mathbb{R}^{n+1}$  joining two points on the graph


**Definition.** A function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is **convex** if

- (i) The domain of  $f$  is a convex set
- (ii)

$$f((1-t)\mathbf{x} + t\mathbf{y}) \leq (1-t)f(\mathbf{x}) + tf(\mathbf{y}) \quad 0 < t < 1 \quad (5.1)$$



convex



non-convex

$f$  is convex if the graph of  $f$  lies below or on its chords

**Remarks.**

- (i)  $f$  is concave if we replace  $\leq$  by  $\geq$  in (5.1)
- (ii)  $f$  convex  $\iff -f$  concave
- (iii)  $f$  strictly convex if we replace  $\leq$  by  $<$  in (5.1)

**Example.**  $f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(x) = x^2$  domain  $\mathbb{R}$  (convex)

$$\begin{aligned} f((1-t)x + ty) - (1-t)f(x) - tf(y) &= [(1-t)x + ty]^2 - (1-t)x^2 - ty^2 \\ &= x^2(1-t) \cdot (-t) + ty^2(1-t) + 2(1-t)txy \\ &= (1-t)t(x-y)^2 < 0 \quad \forall 0 < t < 1 \end{aligned}$$

strictly convex

**Example.**  $f(x) = 1/x$ , domain  $\mathbb{R} \setminus \{0\}$ , not a convex set. On restricted domain  $\mathbb{R} > 0$ ,  $f$  is convex

## 5.1 Conditions for Convexity

3 tests for  $f$  to be convex

(i) If  $f$  is once differentiable, then  $f$  is convex iff

$$f(\mathbf{y}) \geq f(\mathbf{x}) + (\mathbf{y} - \mathbf{x}) \cdot \nabla f(\mathbf{x}) \quad (5.2)$$

**Proof.** Assume (5.2) holds, and apply it twice

$$f(\mathbf{x}) \geq f(\mathbf{z}) + (\mathbf{x} - \mathbf{z}) \cdot \nabla f(\mathbf{z}) \quad (i)$$

$$f(\mathbf{y}) \geq f(\mathbf{z}) + (\mathbf{y} - \mathbf{z}) \cdot \nabla f(\mathbf{z}) \quad (ii)$$

Take  $\mathbf{z} = (1 - t)\mathbf{x} + t\mathbf{y} \in S$  (the domain of  $f$ ),  $0 < t < 1$

$$(1 - t) \cdot (i) + t \cdot (ii) \rightarrow \nabla f(\mathbf{z}) \text{ cancel. get (5.1)}$$

Converse: assume convexity (5.1) and set

$$h(t) = (1 - t)f(\mathbf{x}) + t(f(\mathbf{y})) - f((1 - t)\mathbf{x} + t\mathbf{y}) \geq 0$$

$$h'(0) = -f(\mathbf{x}) + f(\mathbf{y}) - (\mathbf{y} - \mathbf{x}) \cdot \nabla f(\mathbf{x})$$

So (5.2) is equivalent to  $h'(0) \geq 0$ . Note  $h(0) = 0$ , so

$$\frac{h(t) - h(0)}{t} \geq 0 \quad 0 < t < 1$$

Now take the limit  $t \rightarrow 0$

**Corollary.** If  $f$  is convex and have a stationary point, then it is a global minimum

**Proof.** Given  $\nabla f(\mathbf{x}_0) = 0$ , we get from (5.2) that  $f(\mathbf{y}) \geq f(\mathbf{x}_0) \forall \mathbf{y}$

(ii) If

$$(\nabla f(\mathbf{y}) - \nabla f(\mathbf{x})) \cdot (\mathbf{y} - \mathbf{x}) \geq 0 \quad (5.3)$$

then  $f$  is convex ( $f'$  monotonically increasing if  $n = 1$ )

**Proof.** exercise

(iii) (Second order conditions): Assume  $f$  twice differentiable, then  $f$  convex iff the Hessian  $\frac{\partial^2 f}{\partial x^i \partial x^j}$  has all eigenvalues non-negative. If all eigenvalues positive, then  $f$  is strictly convex

**Proof.** Assume convex and apply (5.3) by taking  $\mathbf{y} = \mathbf{x} + \mathbf{h}$

$$\mathbf{h} \cdot (\nabla f(\mathbf{x} + \mathbf{h}) - \nabla f(\mathbf{x})) \geq 0$$

for small  $\mathbf{h}$ :

$$\partial_i f(\mathbf{x} + \mathbf{h}) = \partial_i f(\mathbf{x}) + \sum_j h_j H_{ij}(\mathbf{x}) + O(|\mathbf{h}|^2)$$

So (by dotting with  $\mathbf{h}$ )

$$\sum_{j,i} h_i h_j H_{ij}(\mathbf{x}) + O(|\mathbf{h}|^2) \geq 0$$

**Example.**

$$f(x, y) = \frac{1}{xy} \quad x, y > 0$$

$$H = \frac{1}{xy} \begin{bmatrix} \frac{2}{x^2} & \frac{1}{xy} \\ \frac{1}{xy} & \frac{2}{y^2} \end{bmatrix} \quad \det(H) = \frac{3}{x^3 y^3} > 0 \quad \text{tr}(H) > 0$$

so  $f$  is strictly convex


## 6 Legendre Transform

**Definition.** The Legendre transform of  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is

$$f^*(\mathbf{p}) = \sup_{\mathbf{x}} (\mathbf{p} \cdot \mathbf{x} - f(\mathbf{x})) \tag{6.1}$$

The domain of  $f^*$  consists of all vectors  $\mathbf{p} \in \mathbb{R}^n$  s.t. the sup is finite

**Example.**  $n = 1$



Maximum vertical distance between graphs of  $z = f(x)$  and  $z = px$



**Example.**  $n = 1$ ,  $f(x) = ax^2$ ,  $a > 0$

$$f^*(p) = \sup_x (px - ax^2) \quad \frac{\partial}{\partial x} (px - ax^2) = 0 \implies p = 2xa$$

So  $x = p/2a$  and substitute

$$f^*(p) = p \frac{p}{2a} - a \left(\frac{p}{2a}\right)^2 = \frac{p^2}{4a}$$

Compute  $(f^*)^*(s) = \sup_p (sp - \frac{p^2}{4a}) \implies p = 2as$

$$f^{**}(s) = as^2$$

so  $f^{**} = f$  (always true if  $f$  convex)

If  $a < 0$ ,  $\sup_x (px - ax^2) = \infty \forall p$  so  $f^*$  has empty domain

**Prop.** Domain of  $f^*$  is a convex set, find  $f^*$  convex

**Proof.**

$$f^*((1-t)\mathbf{p} + t\mathbf{q}) = \sup_{\mathbf{x}} [(1-t)\mathbf{p} \cdot \mathbf{x} + t\mathbf{q} \cdot \mathbf{x} - f(\mathbf{x})] = \sup_{\mathbf{x}} [(1-t)[\mathbf{p} \cdot \mathbf{x} - f(\mathbf{x})] + t[\mathbf{q} \cdot \mathbf{x} - f(\mathbf{x})]]$$

Use  $\sup(A+B) \leq \sup(A) + \sup(B)$  to get

$$LHS \leq (1-t)f^*(\mathbf{p}) + tf^*(\mathbf{q})$$

(i)

$$(1-t)\mathbf{p} + t\mathbf{q} \in D(f^*)$$

(ii)  $f^*$  satisfies convexity definition (5.1)

**Note.** In practice, if  $f$  convex and differentiable,

$$f^*(\mathbf{p}) = \text{global minimum over } \mathbf{x}$$

$$\nabla(\mathbf{p} \cdot \mathbf{x} - f(\mathbf{x})) = 0 \implies \mathbf{p} = \nabla f$$

(substitute to definition of  $f^*(p)$ )

If  $f$  is strictly convex, then  $\exists$  unique inversion  $\mathbf{x} = \mathbf{x}(\mathbf{p})$  so that

$$f^*(\mathbf{p}) = \mathbf{p} \cdot \mathbf{x}(\mathbf{p}) - f(\mathbf{x}(\mathbf{p})) \tag{6.2}$$

## 6.1 Applications to Thermodynamics

Many particles (gas  $\sim 10^{23}$  particles) so we use a few macroscopic variables:  $p$  (pressure),  $V$  (volume),  $T$  (temperature),  $S$  (entropy). (Part II Statistical Physics)  
Internal energy  $U(S, V)$ . Helmholtz free is defined

$$F(T, V) = \min_S (U(S, V) - TS) = -\max_S (TS - U(S, V)) = -U^*(T, V)$$

Legendre transform of  $U$  w.r.t.  $S$ , with  $V$  held fixed as a parameter

$$\frac{\partial}{\partial S} (TS - U(S, V))|_{T, V} = 0 \rightarrow T = \frac{\partial U}{\partial S}|_V$$


Other quantities as Legendre transform e.g. ~~Entropy~~ **Enthalpy**

$$H(S, p) = \min_V (U(S, V) + pV) = -U^*(-p, S)$$

at min

$$p = - \left( \frac{\partial U}{\partial V} \right) |_S$$

Entropy is a fixed parameter. The Legendre transform is a way to swap from  $(S, V)$  dependence to dependence of other variables



## 7 Hamilton's Equations

**Remark.** Recall (section 4.1) Lagrangian  $L = T - V = L(\mathbf{q}, \dot{\mathbf{q}}, t)$  function on the configuration space

**Definition.** The Hamiltonian is the Legendre transform of  $L$  w.r.t.  $\dot{\mathbf{q}} = \mathbf{v}$

$$H(\mathbf{q}, \mathbf{p}, t) = \sup_{\mathbf{v}} (\mathbf{p} \cdot \mathbf{v} - L) = \mathbf{p} \cdot \mathbf{v} - L(\mathbf{q}, \mathbf{v}, t)$$

where  $\mathbf{v} = \mathbf{v}(\mathbf{p})$  is the solution to

$$p_i = \frac{\partial L}{\partial \dot{q}_i}$$

(assume convexity of  $L$  in  $\mathbf{v}$ ).  $p$  is the generalised momentum

Example.

$$T = \frac{1}{2}m|\dot{\mathbf{q}}|^2 \quad V = V(\mathbf{q})$$

$$\mathbf{p} = \frac{\partial L}{\partial \dot{\mathbf{q}}} = m\dot{\mathbf{q}} \rightarrow \dot{\mathbf{q}} = \frac{\mathbf{p}}{m}$$

$$\begin{aligned} H(\mathbf{q}, \mathbf{p}, t) &= \mathbf{p} \cdot \frac{\mathbf{p}}{m} - \left( \frac{1}{2}m \frac{|\mathbf{p}|^2}{m^2} - V(\mathbf{q}) \right) \\ &= \frac{1}{2m}|\mathbf{p}|^2 + V(\mathbf{q}) \quad (\text{the total energy}) \end{aligned}$$

What happened to the Euler-Lagrange equations? Assume  $H$  satisfies E-L.

$$H = H(\mathbf{q}, \mathbf{p}, t) = p_i \dot{q}^i - L(q^i, \dot{q}^i, t) \quad \text{Using summation convention}$$

$$dH = \frac{\partial H}{\partial q^i} dq^i + \frac{\partial H}{\partial p_i} dp_i + \frac{\partial H}{\partial t} dt \quad \text{doesn't follow from this.}$$

$$\begin{aligned} dH &= p_i \dot{q}^i + \dot{q}^i dp_i - \frac{\partial L}{\partial q^i} dq^i - \frac{\partial L}{\partial \dot{q}^i} \dot{q}^i - \frac{\partial L}{\partial t} dt \\ &= \dot{q}^i dp_i - \dot{p}_i dq^i - \frac{\partial L}{\partial t} dt \end{aligned}$$

by E-L. Compare differentials

$$\dot{q}^i = \frac{\partial H}{\partial p_i} \quad \dot{p}_i = -\frac{\partial H}{\partial q^i} \quad \frac{\partial H}{\partial t} = -\frac{\partial L}{\partial t} \quad \frac{dL}{dq^i} = \frac{d}{dt} \left( \frac{\partial L}{\partial \dot{q}^i} \right) \text{ by E-L.} = \dot{p}_i \quad (7.2)$$

**Warning.**

$$\frac{\partial}{\partial t} \Big|_{p, q} \neq \frac{\partial}{\partial t} \Big|_{q, \dot{q}}$$

Assume no explicit  $t$ -dependence in  $L$ . Then (7.2) is a system of  $2n$  1st order ODEs. Need to specify  $q^i(0), p_i(0), i = 1, \dots, n$ . Solution curves to (7.2) are trajectories in  $2n$ -dimensional phase space

**Remark.** Hailton's equations also arise from extremizing a functional in phase space

$$S[\mathbf{q}, \mathbf{p}] = \int_{t_1}^{t_2} \underbrace{(\dot{q}^i p_i - H(\mathbf{q}, \mathbf{p}, t))}_{f(\mathbf{q}, \mathbf{p}, \dot{\mathbf{q}}, \dot{\mathbf{p}}, t)} dt$$

E-L for  $S$

- Variation w.r.t.  $p_i$

$$\frac{\partial f}{\partial p_i} - \underbrace{\frac{d}{dt} \left( \frac{\partial f}{\partial \dot{p}_i} \right)}_0 = 0 \implies \dot{q}^i = \frac{\partial H}{\partial p_i}$$

- Variation w.r.t.  $q^i$

$$\frac{\partial f}{\partial q^i} - \frac{d}{dt} \left( \frac{\partial f}{\partial \dot{q}^i} \right) = 0 \implies \cancel{\frac{\partial H}{\partial q^i}} - \frac{\partial H}{\partial q^i} - \frac{dp_i}{dt} = 0$$

$$\dot{p}_i = -\frac{\partial H}{\partial q^i}$$

Recovered (7.2), Newton's equation, Lagrange's equation, Hamilton's equation so far (7.2) is just another formulation

## 8 The Second Variation

E-L equation gives us necessary condition so we could get a minimum, maximum or a saddle point. And so we look at the nature of stationary points of

$$F[y] = \int_{\alpha}^{\beta} f(x, y, y') dx$$

Expand  $F[y + \varepsilon \eta]$  to 2nd order in  $\varepsilon$  around a solution to E-L equation

$$\begin{aligned} F[y + \varepsilon \eta] - F[y] &= \int_{\alpha}^{\beta} [f(x, y + \varepsilon \eta, y' + \varepsilon' \eta') - f] dx \\ &= 0 + \varepsilon \int_{\alpha}^{\beta} \underbrace{\eta \left( \frac{\partial f}{\partial y} - \frac{d}{dx} \left( \frac{\partial f}{\partial y'} \right) \right)}_0 dx + \frac{\varepsilon^2}{2} \int_{\alpha}^{\beta} \left[ \eta^2 \frac{\partial^2 f}{\partial y^2} + (\eta')^2 \frac{\partial^2 f}{\partial (y')^2} + 2 \frac{\partial^2 f}{\partial y \partial y'} \eta \eta' \right] dx \\ &\quad + O(\varepsilon^3) \end{aligned}$$

2nd variation is

$$\begin{aligned} \delta^2 F[y] &\equiv \frac{1}{2} \int_{\alpha}^{\beta} \left[ \eta^2 \frac{\partial^2 f}{\partial y^2} + (\eta')^2 \frac{\partial^2 f}{\partial (y')^2} + \frac{d}{dx} (\eta^2) \frac{\partial^2 f}{\partial y' \partial y} \right] dx \\ &= \frac{1}{2} \int_{\alpha}^{\beta} Q \eta^2 + P (\eta')^2 dx \end{aligned}$$

ibp:  $\int_{\alpha}^{\beta} \eta^2 \frac{d}{dx} \frac{\partial^2 f}{\partial y' \partial y} dx = \int_{\alpha}^{\beta} \eta^2 \frac{d}{dx} \frac{\partial^2 f}{\partial y' \partial y} dx$

where

$$P = \frac{\partial^2 f}{\partial (y')^2} \quad Q = \frac{\partial^2 f}{\partial y^2} - \frac{d}{dx} \left( \frac{\partial^2 f}{\partial y' \partial y} \right) \quad (8.1)$$

we have proved

**Prop.** If  $y(x)$  is a solution to the E-L equation (2.3) and  $Q\eta^2 + P(\eta')^2 > 0 \forall \eta$  vanishing at  $\alpha, \beta$  then  $y(x)$  is a local minimizer of  $F[y]$

**Example.** Geodesics on a plane (in section 2)

$$f = \sqrt{1 + (y')^2} : \begin{cases} P = \frac{\partial}{\partial y'} \left( \frac{y'}{\sqrt{1+(y')^2}} \right) \rightarrow \frac{1}{(1+(y')^2)^{3/2}} > 0 \\ Q = 0 \end{cases}$$

If  $\eta' = 0$ , then  $\eta = 0$ , so  $\eta' \neq 0$  and  $P(\eta')^2 > 0 \forall \eta$  so straight lines are local length minimizers on  $\mathbb{R}^2$


**Prop.** If  $y_0(x)$  is a local minimum, then

$$P = \frac{\partial^2 f}{\partial (y')^2} \Big|_{y_0} \geq 0 \tag{8.2}$$

so the Legendre condition is necessary for local min.

“ $P$  is more important than  $Q$  in (8.1)”

**Proof.** See Gelfand-Fomin for details. Idea: if  $\eta'$  small, then  $\eta$  can't be too large. Converse not true:  $\eta$  can be small,  $\eta'$  large. Assume  $\exists x_0$  s.t.  $P(x_0, y_0, y'_0) < 0$



**Note.** (8.2) not sufficient for local minimum see section 8.1 but  $P > 0, Q \geq 0$  is sufficient as if  $\eta \neq 0$  on  $(\alpha, \beta)$  then  $\exists x_0 \in (\alpha, \beta)$  s.t.  $\eta'(x_0) \neq 0$

**Example.** Go back to Brachistochrome

$$f = \sqrt{\frac{1 + (y')^2}{-y}}$$

Is cycloid a minimizer?

$$\frac{\partial f}{\partial y} = -\frac{1}{2y} f \quad \frac{\partial f}{\partial y'} = \frac{y'}{\sqrt{1 + (y')^2} \sqrt{-y}}$$

$$P = \frac{1}{(1 + (y')^2)^{3/2} \sqrt{-y}} > 0$$

$$Q = \dots = \frac{1}{2\sqrt{1 + (y')^2} y^2 \sqrt{-y}}$$

## 8.1 Associated Eigenvalue Problem

Go back to (8.1)

$$Q\eta^2 + P(\eta')^2 = Q\eta^2 + \frac{d}{dx}(P\eta\eta') - \eta(P\eta)'$$

$\int_{\alpha}^{\beta} P\eta'^2 = \left[ \eta P\eta' \right]_{\alpha}^{\beta} - \int_{\alpha}^{\beta} \eta (P\eta)'$

integrate, drop the boundary term as  $\eta = 0$  at  $\alpha, \beta$

$$\delta^2 F[y_0] = \frac{1}{2} \int_{\alpha}^{\beta} \underbrace{\eta[-(P\eta)'] + Q\eta}_{\mathcal{L}(\eta)} dx \quad (8.3)$$

Sturn-Liouville operator. If  $\exists \eta$  s.t.

$$\begin{cases} \mathcal{L}(\eta) = -\omega^2 \eta \quad (\omega \text{ real}) \\ \eta(\alpha) = \eta(\beta) = 0 \end{cases} \quad (8.4)$$

Then  $y_0$  is not a minimizer as

$$\delta^2 F[y_0] = -\frac{1}{2} \omega^2 \int_{\alpha}^{\beta} \eta^2 dx < 0$$

(8.4) can have solutions even if  $P > 0$ , so the Legendre condition (8.2) is not sufficient for  $y_0$  to be a minimizer

**Example.**

$$F[y] = \int_0^{\beta} [(y')^2 - y^2] dx$$

with  $y(0) = y(\beta) = 0$  and  $\beta \neq N\pi$   $N \in \mathbb{N}$

$$(2.3) \rightarrow y'' + y = 0 \implies y = y_0 = 0$$

is the stationary point of  $F[y]$ . 2nd variation:

$$\delta^2 F[0] = \frac{1}{2} \int_0^{\beta} [(\eta')^2 - \eta^2] dx \quad P = \frac{2}{4} > 0$$

but  $Q < 0$ . Examine (8.4):

$$-\eta'' - \eta = -\omega^2 \eta \quad \eta(0) = \eta(\beta) = 0$$

Take

$$\eta = A \cdot \sin\left(\frac{\pi x}{\beta}\right) \rightarrow \left(\frac{\pi}{\beta}\right)^2 = 1 - \omega^2$$

Possible if  $\beta > \pi$ . So, if  $P > 0$  a problem may arise if the interval is "too large".

## 8.2 The Jacobi Condition

Legendre tried to prove that  $P > 0$  is sufficient for  $y = y_0$  to be a local minimum. This couldn't have worked (last example), but the idea was good.

Let  $\phi = \phi(x)$  be a any differentiable function of  $x$  on  $[\alpha, \beta]$

$$0 = \int_{\alpha}^{\beta} (\phi\eta^2)' dx = \int_{\alpha}^{\beta} \phi'\eta^2 + 2\eta\eta'\phi dx \quad \left[ \phi\eta^2 \right]_{\alpha}^{\beta}$$

(as  $\eta(\alpha) = \eta(\beta) = 0$ ). Adding to (8.1), we can rewrite

$$0 + \delta^2 F[y] = \frac{1}{2} \int_{\alpha}^{\beta} (P(\eta')^2 + 2\eta\eta'\phi + (Q + \phi')\eta^2) dx$$

Assume  $P|_y > 0$  and complete the square

$$\delta^2 F[y] = \frac{1}{2} \int_{\alpha}^{\beta} \left[ P\left(\eta' + \frac{\phi}{P}\eta\right)^2 + \underbrace{\left(Q + \phi' - \frac{\phi^2}{P}\right)}_{=0 \text{ if (8.3) holds}} \eta^2 \right] dx$$

which is positive if we can choose  $\phi$  s.t.

$$\phi^2 = P(Q + \phi') \quad (8.3)$$

If (8.3) holds, then  $\delta^2 F > 0$  unless

$$\eta' + \frac{\phi}{P}\eta = 0 \quad (**)$$

on  $[\alpha, \beta]$ . But  $\eta = 0$  at  $\alpha$ , so  $\eta'(\alpha) = 0$  if (\*\*) holds but then  $\eta \equiv 0$  on  $[\alpha, \beta]$  (uniqueness of solution to 1st order ODEs), so (\*\*)  $\neq 0$ .

**Method.** Does a solution to (8.3) exist on  $[\alpha, \beta]$ ?

Transform (8.3) into a linear 2nd order ODE by setting  $\phi = -Pu'/u$  where  $u \neq 0$  on  $[\alpha, \beta]$

$$P\left(\frac{u'}{u}\right)^2 = Q - \left(\frac{Pu'}{u}\right)' = Q - \frac{(Pu)'}{u} + P\left(\frac{u'}{u}\right)^2$$

or

$$-(Pu')' + Qu = 0 \quad (8.4)$$

This is the Jacobi accessory condition.

Need a solution to (8.4) (which is  $\mathcal{L}(u) = 0$ ) s.t.  $u \neq 0$  on  $[\alpha, \beta]$ . This may not exist on a large enough interval

**Example.**

$$F[y] = \frac{1}{2} \int_{\alpha}^{\beta} [(y')^2 - (y^2)] dx$$

$$y \rightarrow y + \varepsilon \eta \quad \delta^2 F[y] = \frac{1}{2} \int_{\alpha}^{\beta} [(\eta')^2 - \eta^2] dx \quad P = 1, Q = -1$$

(8.4) is  $u'' + u = 0$ , general solution  $u = A \sin x + B \cos x$ . Want  $u$  to be non-zero on  $[\alpha, \beta]$ , i.e.

$$\tan(x) \neq \frac{B}{A}$$

possible to avoid  $B/A$  on interval smaller than  $\pi$  2<sup>nd</sup>

$$|\beta - \alpha| < \pi \rightarrow \text{positive } \eta \text{ variation}$$

**Example.** Back to geodesics on the sphere

$$\delta \sqrt{d\theta^2 + \sin^2 \theta d\phi^2} = \sqrt{(\theta')^2 \sin^2 \theta d\theta^2} \quad \theta = \theta(\phi)$$

f

Found earlier that critical points are segments of great circles  
 $\theta = \text{const}$ ,  $\theta_0 = \pi/2$  (any great circle is this after a rotation)

$$\frac{\partial^2 f}{\partial (\theta')^2} |_{\theta_0} = 1 = P \quad Q = \dots = -1$$

$$\delta^2 F[\theta_0 = \frac{\pi}{2}] = \frac{1}{2} \int_{\phi_1}^{\phi_2} [(\eta')^2 - \eta^2] d\phi$$

1<sup>st</sup>

positive if  $\phi_2 - \phi_1 < \pi$